

On the distribution of high-frequency stock market traded volume: a dynamical scenario

Silvio M. Duarte Queirós
*Centro Brasileiro de Pesquisas Físicas, 150,
 22290-180, Rio de Janeiro - RJ, Brazil*
Electronic mail address: sdqueiro@cbpf.br

(Dated: February 2, 2008)

This manuscript reports a stochastic dynamical scenario whose associated stationary probability density function is exactly a previously proposed one to adjust high-frequency traded volume distributions. This dynamical conjecture, physically connected to superstatistics, which is intimately related with the current nonextensive statistical mechanics framework, is based on the idea of local fluctuations in the mean traded volume associated to financial markets agents herding behaviour. The corroboration of this mesoscopic model is done by modelising NASDAQ 1 and 2 minute stock market traded volume.

PACS numbers: 05.90.+m, 05.40.-a, 89.65.Gh.

Keywords: financial markets; traded volume; nonextensive statistical mechanics

In the last years we have assisted to a rapidly growing interesting of physicists in the analysis of financial systems and the description of their statistical and dynamical properties [1, 2, 3, 4, 5]. Actually, some of the so-called *stylised facts* such as non-Gaussianity in relative price change (the *return*) probability density functions (PDFs) [6, 7, 8, 9, 10, 11], multi-fractal behaviour of return time series [12, 13, 14], asymptotic power-law like auto-correlation function in the standard deviation (the *volatility*) of return [15, 16], among many others, were (also) verified in a physical context. Besides their own relevance, these empirical observations were important to the establishment of either new theoretical models, which mimic reality much better than previous ones [16, 17, 18, 19], or connections between financial systems and already existing statistical frameworks [10, 20, 21], particularly, the current non-extensive statistical mechanics [5], based on *Tsallis entropy* [22], which, *e.g.*, allowed the derivation of a new and closed model for option-pricing [23, 24, 25].

Additionally to macroscopic observables as the return and the volatility, mentioned above, another significant quantity in financial markets is the *traded volume*, V , which corresponds to the number of shares traded (in a determined period of time, T). This is, in fact, along with volatility, an important parameter in trading strategies assumed by many brokers [26, 27]. In fig. 1 is presented, as mere illustration, the daily normalised traded volume of Microsoft Corp. in NASDAQ exchange from 2000 to the beginning of 2005.

In a previous work, P. Gopikrishnan *et al.* [28] reported the evidence, for large volumes, of a power-law behaviour in the stationary PDF. The analysis of traded volume stationary PDF was later extended to all range of values by R. Osorio, L. Borland and C. Tsallis (OBT) [10], who proposed a fitting expression that presents an amazing precision. The respective formula is,

$$p(v) = \frac{1}{Z} \left(\frac{v}{\theta} \right)^\alpha \exp_q \left(-\frac{v}{\theta} \right), \quad (1)$$

where v represents the traded volume expressed in its mean value unit ($\langle V \rangle$), *i.e.*, $v_t = \frac{V_t}{\langle V \rangle}$, $Z = \int_0^\infty \left(\frac{v}{\theta} \right)^\alpha \exp_q \left(-\frac{v}{\theta} \right) dv$, α and θ are positive parameters and

$$\exp_q(x) \equiv [1 + (1 - q)x]^{\frac{1}{1-q}}, \quad (\exp_1(x) = e^x), \quad (2)$$

is the *q-exponential function*, which emerged in the context of the *non-extensive statistical mechanics* [5].

In the sequel of this article, through a set of liable arguments, we will introduce a stochastic dynamical mechanism, which has eq. (1) as stationary solution for its probability density function and then compare both analytical and simulation results using values obtained from NASDAQ stock market.

Consider the following stochastic differential equation,

$$dv = -\gamma \left(v - \frac{\alpha + 1}{\beta} \right) dt + \sqrt{2v \frac{\gamma}{\beta}} dW_t, \quad (3)$$




FIG. 1: Evolution of Microsoft Corporation daily normalised traded volume in NASDAQ exchange from the 21st January 2001 to the 21st January 2005. Each tick represents 6 months.

(W_t represents a regular Wiener process following a normal distribution with null mean and unitary variance) as the traded volume updating law ¹. As can be seen, eq. (3) is composed by a deterministic term, similar to a restoring force and a stochastic term, which is a typical example of multiplicative noise very usual in problems presenting clustering profiles, like the one treated in this manuscript (see fig. 1), intermittence [30] or turbulence [31, 32]. The first term can be interpreted as a restoring market mechanism (with constant γ), which aims to keep the traded volume of the stock at some typical value $\Theta = \frac{\alpha+1}{\beta}$. ² The second one reflects stochastic memory, and mainly, the upshot of large traded volumes. In effect, large traded volumes will induce large values in the stochastic term, but, due to W_t random character(sign), it could lead to an increase ($W_t > 0$) or a decrease ($W_t < 0$) of v , *i.e.*, to a stirred or serene state of the stock.

For eq. (3) we can write the corresponding Fokker-Planck equation [33],

$$\frac{\partial p(v,t)}{\partial t} = \frac{\partial}{\partial v} \left[\gamma \left(v - \frac{\alpha+1}{\beta} \right) p(v,t) \right] + \frac{\partial^2}{\partial v^2} \left[v \frac{\gamma}{\beta} p(v,t) \right], \quad (4)$$

and determining its stationary solution we obtain,

$$p(v) = \frac{\beta}{\Gamma[1+\alpha]} (\beta v)^\alpha \exp(-\beta v) \quad (\alpha > -1, \beta > 0), \quad (5)$$

which is just the Gamma distribution, with mean value, $\langle v \rangle = \frac{1+\alpha}{\beta}$, standard deviation, $\langle v - \langle v \rangle \rangle^2 = \frac{1+\alpha}{\beta^2}$ and most probable value, $v^* = \frac{\alpha}{\beta}$ ($\frac{dP(v)}{dv} \Big|_{v=v^*} = 0$).

Assume now that, instead of constant in time, β is now randomly varying on a time scale much larger than the time scale of order γ^{-1} necessary by eq. (4) to reach stationarity, hypothesis that is equivalent to consider local fluctuations in the mean value of v [34], akin to the proposal of moving average in the volatility [35]. These fluctuations can be microscopically associated to changes in the *volume of activity* defined as the number of traders who performed transactions of the stock at that time [36]. Furthermore, let us assume that β follows a Gamma (stationary) PDF,

$$P(\beta) = \frac{1}{\lambda \Gamma[\delta]} \left(\frac{\beta}{\lambda} \right)^{\delta-1} \exp\left(-\frac{\beta}{\lambda}\right) \quad (\delta > 0, \lambda > 0). \quad (6)$$

This assumption transforms eq. (5) meaning from the absolute probability of v into the conditional probability of v given β ,

$$p(v) \rightarrow p(v|\beta) = \frac{\beta}{\Gamma[1+\alpha]} (\beta v)^\alpha \exp(-\beta v). \quad (7)$$

¹ Eq. (3) is in fact defined in mathematics as *Feller process* [29]. According to its definition the traded volume V is an integer quantity. However, the normalised traded volume is a rational quantity and so perfectly compatible with a continuous approach.

² Following v definition, Θ should equals 1. Nonetheless, we will keep it in the presented form for reasons that will become clear later on.

It is noteworthy that the problem of macroscopic non-equilibrium systems with complex dynamics in stationary states with fluctuations of intensive quantities on long time scales and its connection with non-extensive statistical mechanics, based on entropy [22],

$$S_q = \frac{1 - \sum_i [p_i]^q}{q - 1},$$

($S_1 = -\sum_i p_i \ln p_i \equiv S_{BG}$ where BG stands for Boltzmann-Gibbs) was firstly presented by G. Wilk and Z. Włodarczyk [37], later applied by C. Beck to the case of turbulent fluids [38] and finally extended and defined by C. Beck and E.G.D. Cohen as *superstatistics* [39, 40].

For this case, we have that the *joint* probability of obtaining a certain value of v and certain value of β , is $P(v, \beta) = p(v|\beta) P(\beta)$ and the *marginal* probability of having some value v is given by

$$P(v) = \int_0^\infty P(v, \beta) d\beta = \int_0^\infty p(v|\beta) P(\beta) d\beta. \quad (8)$$

Using eq. (6) and eq. (7) in eq. (8) and performing the integration we get,

$$P(v) = \frac{\lambda \Gamma[1 + \alpha + \delta]}{\Gamma[1 + \alpha] \Gamma[\delta]} (\lambda v)^\alpha (1 + \lambda v)^{-1 - \alpha - \delta} \quad (\alpha + \delta > -1). \quad (9)$$

Doing the change in parameters

$$\lambda = \frac{q - 1}{\theta}, \quad \delta = \frac{1}{q - 1} - \alpha - 1, \quad (10)$$

one can write eq. (9) as

$$P(v) = \frac{(q - 1)^{\alpha + 1} \Gamma\left[\frac{1}{q - 1}\right]}{\theta \Gamma\left[\frac{1}{q - 1} - \alpha - 1\right] \Gamma[\alpha + 1]} \left(\frac{v}{\theta}\right)^\alpha \left[1 + (q - 1) \frac{v}{\theta}\right]^{1/(1 - q)},$$

or, using the q -exponential function definition, eq. (2),

$$P(v) \equiv \frac{(q - 1)^{\alpha + 1} \Gamma\left[\frac{1}{q - 1}\right]}{\theta \Gamma\left[\frac{1}{q - 1} - \alpha - 1\right] \Gamma[\alpha + 1]} \left(\frac{v}{\theta}\right)^\alpha \exp_q\left(-\frac{v}{\theta}\right), \quad (11)$$

which is *exactly* the expression introduced by OBT. Due to its structure, PDF (11) might be called *q -generalised Gamma probability density function*. The traditional Gamma PDF, eq. (5), can be recovered from eq. (11) by considering $q = 1$ limit, which is equivalent to assume the inexistence of fluctuations in β and so $P(\beta)$ equals the Dirac delta function centred in θ^{-1} . For small values of v , $P(v)$ goes to zero as a power law, $P(v) \sim v^\alpha$, and for large values of v , it behaves asymptotically as another power law, $P(v) \sim v^{\alpha/(1 - q)}$. As a matter of fact, this is an appealing form, since it congregates, in a *single expression*, the two power laws used in financial practice.

In order to verify the correctness of the result, we performed a set of numerical realisations, using an algorithm based on eq. (3) and on a random generator for β whose associated probability is eq. (6). The values of parameters α , λ and δ , presented in table I, were obtained after fitting the 1 minute and 2 minute traded volume probability density functions of the ten high-volume stocks in the NASDAQ exchange during 2001, fig. 2, with eq. (1) (equivalent to eq. (11)). The values of q and θ are obtained using the relations presented in eq. (10).

The numerical realisations for 1 minute and 2 minute traded volume are depicted in fig. 3 and fig. 4.

In fig. 1, fig. 3 and fig. 4 (Centre plots) it is clear the clustering behaviour, very similar to volatility time series, typical of a multiplicative noise dynamics. In fig. 3 and fig. 4 (Right plots), we can compare numerically obtained relative frequency, $H(v)$, with analytical results corresponding to values presented in table I. As it is straightforwardly visible, numerical and analytical results are in perfect accordance. Besides the possibility of mimic traded volume time series, this dynamical conjecture can certainly be used in other problems with time varying positive quantities associated to a PDF (1), such as granular material systems [41, 42].

To summarise, in this manuscript we presented a dynamical conjecture, based on a multiplicative noise mechanism and on the fluctuating character of some parameter of that system, which is able to provide the stationary probability




FIG. 2: 1 minute (Left plot) and 2 minute (Right plot) traded volume probability density function $P(v)$, for the ten high-volume stocks in NASDAQ exchange during 2001 *vs.* v . The symbols represent real data and lines $p(v)$ from eq.(1), with q, α, θ values shown in table I (After OBT in ref.[10]). The χ^2 (per point) error function values are 1.8×10^{-3} and 6.5×10^{-4} , respectively. For very low volumes is visible a small discrepancy, due to lack of small transactions (statistical effect).

TABLE I: Fitting parameters obtained in fig.2 and dynamical parameters obtained through eq.10.

	q	α	θ	λ	δ
1 min	1.19	0.93	0.23	0.826	3.33
2 min	1.16	1.36	0.2	0.8	3.89

density function previously introduced to shape the high-frequency traded volume stationary probability distribution of shares in stock markets. The present scenario, related with superstatistics and thus with non-extensive statistical mechanics, contains two ingredients that are often considered as elemental in the dynamics of such complex systems: memory (via multiplicative noise) and microscopic mechanisms of herding between traders (through the local temporal fluctuations of β or the v mean value) caused, *e.g.*, by news, rumours or simply price movements. The description of a completely related return-volatility-volume mesoscopic dynamical theory, within non-extensive statistical mechanics framework, for financial markets is currently in progress.

SMDQ would like to thank to C. Tsallis for his continuous incentive and relevant conversations and suggestions; E.P. Borges is acknowledged for preliminary discussions about PDF form (1) possible origins and comments made on previous versions of this manuscript as well as L.G. Moyano. This work was developed with infrastructural support




FIG. 3: Numerical realisation for 1 minute traded volume of the ten high-volume stocks in NASDAQ exchange during 2001 using the values of table I. Left Plot: temporal evolution of β parameter; Centre Plot: temporal evolution of traded volume, v . Right Plot: Relative frequency $H(v)$ *vs.* v . The symbols were obtained numerically and the line from $\int_{v-\Delta v}^{v+\Delta v} P(v) dv$ with $\Delta v = 5 \times 10^{-3}$. The ratio between β updating time and stationarity time scale of v is 10^2 .




FIG. 4: Numerical realisation for 2 minute traded volume of the ten high-volume stocks in NASDAQ exchange during 2001 using the values of table I. Left Plot: temporal evolution of β parameter; Centre Plot: temporal evolution of traded volume, v . Right Plot: Relative frequency $H(v)$ vs. v . The symbols were obtained numerically and the line from $\int_{v-\Delta v}^{v+\Delta v} P(v) dv$ with $\Delta v = 5 \times 10^{-3}$. The ratio between β updating time and stationarity time scale of v is 10^2 .

from CNPq and PRONEX/MCT (Brazilian agencies) and financial support from FCT/MCES (Portuguese agency; contract SFRH/BD/6127/2001).

-
- [1] J.P. Bouchaud and M. Potters, *Theory of Financial Risks: From Statistical Physics to Risk Management*, (Cambridge University Press, Cambridge, 2000).
 - [2] R.N. Mantegna and H.E. Stanley, *An introduction to Econophysics: Correlations and Complexity in Finance*, (Cambridge University Press, Cambridge, 1999).
 - [3] J. Doyne Farmer, Int. J. Appl. Finance **3**, 311 (2000).
 - [4] J. Voit, *The Statistical Mechanics of Financial Markets (Texts and Monographs in Physics)*, (Springer-Verlag, Berlin, 2003).
 - [5] M. Gell-Mann and C. Tsallis (eds.), *Nonextensive Entropy - Interdisciplinary Applications*, (Oxford University Press, New York, 2004).
 - [6] R.N. Mantegna and H.E. Stanley, *Nature* (London) **376**, 46 (1995).
 - [7] V. Plerou, P. Gopikrishnan, L.A. Amaral, M. Meyer and H.E. Stanley, Phys. Rev. E **60**, 6519 (1999).
 - [8] F.M. Ramos, C. Rodrigues Neto, R.R. Rosa, L.D. Abreu Sá and M.J.A. Bolzam, Nonlinear Analysis **23**, 3521 (2001).
 - [9] M. Ausloos and K. Ivanova, Phys. Rev. E **68**, 046122 (2003).
 - [10] R. Osorio, L. Borland and C. Tsallis in *Nonextensive Entropy - Interdisciplinary Applications*, M. Gell-Mann and C. Tsallis (eds.) (Oxford University Press, New York, 2004).
 - [11] S. M. Duarte Queirós, Physica A **344**, 279 (2004).
 - [12] S. Ghashghaie, W. Breymann, J. Peinke, P. Talkner and Y. Dodge, *Nature* **381**, 767 (1996).
 - [13] J.P. Bouchaud, M. Potters and M. Meyer, Eur. Phys. J. B **13**, 595 (1999).
 - [14] A. Arnéodo, J.-F. Muzy and D. Sornette, Eur. Phys. J. B **2**, 277 (1998).
 - [15] Z. Ding, C.W.J. Granger and R.F. Engle, J. Empirical Finance **1**, 83 (1993).
 - [16] M. Potters, R. Cont and J.-P. Bouchaud, Europhys. Lett. **41**, 239 (1998).
 - [17] L. Borland, Phys. Rev. Lett. **89**, 098701 (2002).
 - [18] M.G. Daniels, J.Doyne Farmer, L. Gillemot, G. Iori and E. Smith, Phys. Rev. Lett. **90**, 108102 (2002).
 - [19] J. Doyne Farmer, L. Gillemot, F. Lillo, S. Mike and A. Sen, Quantitative Finance **4**, 383 (2004).
 - [20] L. Borland, Phys. Rev. Lett. **89**, 098701 (2002).
 - [21] S.M. Duarte Queirós, submitted to Quantitative Finance, 2004.
 - [22] C. Tsallis, J. Stat. Phys. **52**, 479 (1988). A regularly an updated bibliograhly on the subject is available at <http://tsallis.cat.cbpf.br/biblio.htm>.
 - [23] L. Borland, Quantitative Finance **2**, 415 (2002).
 - [24] L. Borland and J.P. Bouchaud, cond-mat/0403022 Preprint, 2004.
 - [25] L. Borland, cond-mat/0412526 Preprint, 2004.
 - [26] P. Willmott, *Paul Willmott on Quantitative Finance*, (Wiley, Chichester, 2000).
 - [27] M. Ausloos and K. Ivanova, in *Application of Econophysics: Proceedings of the Second Nikkei Econophysics Symposium*, H. Takayasu (ed.) (Springer-Verlag, Berlin, 2003).
 - [28] P. Gopikrishnan, V. Plerou, X. Gabaix and H.E. Stanley, Phys. Rev. E **62**, R4493 (2000).
 - [29] W. Feller, Ann. of Math. **54**, 173 (1951).
 - [30] H. Takayasu, A.-H. Sato and M. Takayasu, Phys. Rev. Lett. **79**, 966 (1997).

- [31] U. Frisch, *Turbulence: The Legacy of A. Kolmogorov*, (Cambridge University Press, Cambridge, 1997).
- [32] C. Beck, G.S. Lewis and H.L. Swinney, Phys. Rev. E **63**, R035303 (2001).
- [33] H. Risken, *The Fokker-Planck Equation: Methods of Solution and Applications*, 2nd edition (Springer-Verlag, Berlin, 1989).
- [34] S.M. Duarte Queirós, unpublished working paper available at <http://br.arxiv.org/abs/physics/0411111>, 2004.
- [35] J. Perelló, J. Masoliver and J.P. Bouchaud, Appl. Math. Fin. **11**, 24 (2004).
- [36] T. Lux and M. Marchesi, Nature **397**, 498 (1999).
- [37] G. Wilk, Z. Włodarczyk, Phys. Rev. Lett. **84**, 2770 (2000).
- [38] C. Beck, Phys. Rev. Lett. **87**, 180601 (2001).
- [39] C. Beck and E.G.D. Cohen, Physica A **322**, 267 (2003).
- [40] C. Tsallis and A.M.C Souza, Phys. Rev. E **67**, 026106 (2003).
- [41] E. Kolb, T. Mazozi, E. Clément and J. Duran, Eur. Phys. J. B **8**, 433 1999.
- [42] E.P. Borges, *Manifestações dinâmicas e termodinâmicas de mecânica estatística não-extensiva* (Doctor Thesis), (CBPF, Rio de Janeiro) 2004.